

Sorting Through the Information on Sheep and Goat Parasite Control: A Decision Making Support Tool

Ohio State University Extension

Dr. William Shulaw, Extension Veterinarian, Cattle/Sheep

Rory Lewandowski, Extension Educator, Ag/NR

Dr. Jeff McCutcheon, Extension Educator, Ag/NR

Dr. Jeffrey Workman, Extension Associate, VPM



Copyright © 2012

Version: 1.0



United States Department of Agriculture
National Institute of Food and Agriculture



A Note on Navigation



This tool contains hyperlinks ([text in blue and underlined](#)) that **must** be used for navigation within the tool or to an external website for additional information. There is also an action button located in the upper right corner to return “Home” as well as other buttons to help navigate throughout:



Home - Return to the title screen



Next - Advance to continuation screen



End of Section – Directions to return to table of contents or previous decision.



Please note the following:

- **Caution:** *do not* use the scrolling feature or navigation buttons within your [Adobe Reader](#) toolbar – hyperlinks (including hyperlinked buttons) **must** be used for proper and accurate navigation.
- **Optional feature:** right click your mouse and select “**Previous View**” to return to the last screen viewed.
- **Contact information:** please contact workman.45@osu.edu if you find any hyperlinks that no longer work.

To continue navigation, select “Next” button





DECISION MAKING SUPPORT TOOL


This decision making support tool is designed to help sheep and goat producers sort through the large amount of information available on controlling sheep and goat parasites and make decisions about specific management options that are relevant to their farm operation. It is not intended to replace your veterinarian with regard to diagnosis of parasitism or specifics of drug use. This information is organized in a “decision tree” or “flow chart” approach where answering one question leads to another question or various management options. Each section of the flow chart is basically organized in the format of Class of animal → Time of year → Degree of management flexibility (or availability of resources at the farm level). In some cases, the user will be referred to external references or resources providing additional information on selected topics (e.g., the FAMACHA© system or using certain plants in control of parasitism).



Introduction



DECISION MAKING SUPPORT TOOL – OVERALL LAYOUT



This diagram is an illustration of the overall layout of the decision support tool. The general organization is Class of animal -> Time of year -> Degree of flexibility or available resources at the farm level.

To continue navigation, select "Next" button



Table of Contents



Click on the blue colored text to move to the appropriate section of interest.

[Determining Dewormer Effectiveness](#) – Because some reliance on chemical dewormers is necessary for most farms, it is very important to know how well these products perform against the worms on your farm. Research has clearly shown that using dewormers which are not optimally effective can result in lost productivity and lowered animal welfare. This chart helps you to learn how to determine which dewormers might be effective.

Which class of animals do you wish to learn about?

[Lactating females](#) – lactating females with nursing lambs and kids

[Weaned, growing lambs and kids](#) – young males and females in their first few months of life

[Replacement females](#) – young females, older than 6 months, that will enter the breeding groups

[Dry ewes and does](#) – non-lactating females

[Breeding males](#) – rams and bucks of breeding age and used as flock/herd sires

[Glossary](#) – Definitions to help you understand terms used in this Decision Making Support Tool for managing internal parasites.

[Additional resources](#) – Selected fact sheets, articles, and websites which provide you with additional information about topics discussed in this decision making support tool.

Glossary



Definitions to Help You Understand Terms Used in this Decision Making Support Tool for Managing Internal Parasites

- [Egg-to-larval development](#)
- [DrenchRite® Assay](#)
- [FAMACHA©](#)
- [Fecal egg count reduction testing](#)
- [GIN – “gastrointestinal nematodes”](#)
- [*Haemonchus contortus*](#)
- [Hypobiosis](#)
- [Leader-follower](#)
- [Life cycle](#)
- [Non-persistent dewormer](#)
- [Refugia](#)
- [Rotational grazing](#)
- [Safe Pasture](#)
- [Set stocking](#)
- [Stocking density](#)
- [Strip grazing](#)
- [Summer annual](#)
- [TST – “Targeted Selective Treatment”](#)
- [Winter annual](#)



Egg-to-larval development – Adult worms expel eggs that pass outward in feces (manure). These eggs hatch into larvae on pastures under favorable conditions of moisture and temperature. When the worm larvae are ingested by sheep and goats, they develop into adult worms in the gastrointestinal tract and begin the cycle all over again.

- View short video clip showing [development of worm eggs to third stage larvae](#).
(video created by Dr. Cliff Monahan, OSU Veterinary Parasitologist)

In this video you can watch through the microscope as worm eggs undergo changes toward hatching into the first stage larva and then to the final, or third stage, larva which is infectious for sheep and goats. The third stage larva has a rough or “corrugated-like” appearance to its outer surface. The process of egg development to infective third stage larva can occur in as little as 4 days under ideal conditions. More commonly it takes about 7 days.


Glossary



DrenchRite® Assay – uses eggs harvested from manure samples that are representative of your sheep and goats. The larvae developing from these eggs are exposed to differing levels of dewormers and parasitologists can determine which may be effective. All three chemical classes of dewormers can be tested at one time.

- View short video clip showing [close-up view of three individual wells of the DrenchRite Assay® plate](#). (video created by Dr. Cliff Monahan, OSU Veterinary Parasitologist)

In this video, you can watch as the microscope zooms in on the contents of three individual wells of the DrenchRite® Assay plate. You will see worm larvae that have developed from eggs and a few eggs that did not develop. In this assay, a known number of worm eggs are placed in each of the 96 wells, or cavities, of the plastic plate. Each well contains some nutrients and moisture to support the development of worm larvae to the infectious third stage (L3). The wells contain varying concentrations of dewormers, and the number of larvae developing to the third stage is compared to the number developing in control wells that have no dewormer. In this way, resistance to the three classes of dewormer can be detected.



- Consult the [laboratory](#) at the University of Georgia for DrenchRite® Assay.

Glossary



FAMACHA® – an acronym formed from FAffa MALan CHArt – a system named in honor of one of its South African developers, Dr Francois “Faffa” Malan. This system uses a patented color chart against which the color of the inner surface of the lower eyelid is compared. The colors are 5 shades of red varying from red to very pale pink or flesh color, and they correspond to levels of circulating red blood cells. Bright red is correlated with normal red cell levels and the paler colors correlate with anemia or shortages of red blood cells. The GIN, *Haemonchus contortus*, is found in the abomasum of sheep and goats and feeds on blood. Large numbers of this worm cause anemia, poor performance, and even death. The FAMACHA® system is one method of targeted selective treatment and is applicable to *Haemonchus contortus* only. This is the most serious worm for sheep and goat producers across much of the United States.



More information about the FAMACHA® system can be found at:

<http://www.sheepandgoat.com/ACSRPC/Resources/famacha.html>


FAMACHA® cards and training may be available from your veterinarian or call your Extension educator to request training in your area.

Glossary



Fecal egg count reduction testing – A fecal worm egg count (FEC) is done on manure to look for worm eggs. It is quantitative versus qualitative in that the result is expressed as the number of eggs per gram (epg) of manure as opposed to “positive” or “negative” or “+, ++, or +++” results that are often given from simple flotation procedures. A quantitative result gives us a means to quantify changes over time or in response to a treatment. The three main uses of FECs are to detect dewormer resistance, to monitor pasture contamination, and to select animals for their genetic ability to resist worms.

- See OSUE fact sheet VME-27-11 [“What Do Fecal Worm Egg Counts Tell Us?”](#)
- Consult your veterinarian for fecal egg count reduction testing.




[McMaster slide](#)

Glossary



GIN – “gastrointestinal nematodes” – this is a common abbreviation referring to the roundworm parasites of sheep and goats. It does not refer to tapeworms or coccidia which are not covered in this decision support tool and which do not need pasture for transmission. The important GIN of sheep and goats have similar life cycles which require development of larval stages on pastures. It is used in this decision support tool to save space.



The worms above are images of the important roundworms of sheep and goats (GIN). These worms will not be found in the manure, but their eggs will.



In the image above, the white shapes are segments of the common tapeworm of sheep and goats in clumps of manure.



Haemonchus contortus – or the “barber pole worm” is a blood sucking internal parasite with an extremely high reproductive rate, a quick life cycle, and an ability to survive on pasture paddocks for a long period of time. Under ideal conditions of moderate to high temperatures and moisture, the entire life cycle can be completed in as little as 23 days. The adult *Haemonchus contortus* parasite can lay up to 5,000 eggs per day, which in a grazing operation are shed on to the pasture where the sheep or goats are grazing.

- View short video clip showing [Haemonchus worms collected from the abomasum of a sheep dying of blood loss](#). (video created by Dr. Cliff Monahan, OSU Veterinary Parasitologist)

This video shows a tangled mass of adult Haemonchus contortus worms as found in the abomasal (stomach) contents of a sheep that had died of blood loss created by massive infection with these worms. Enormous numbers of these worms were found in the stomach contents. A blood clot helps hold this mass of worms together. As the tangled mass is teased apart, you can see the individual worms and their “barber pole” appearance caused by the worm’s blood-filled gut spiraling around the large uterus.

Glossary




Hypobiosis – In the normal life cycle, worm eggs passed in the feces develop into larvae that are ingested by the sheep, and the larvae develop into adult worms in the sheep's abomasum or intestines. Sometimes the larvae do not complete their development but go into a resting phase in the abomasal or intestinal tissue. They do relatively little harm in this phase which is called hypobiosis. Day length changes or the onset of milk production in pregnant females stimulates them to continue development to the adult stage. This is one strategy for worms to survive harsh environmental conditions such as northern winters or very hot dry summers.

Glossary



Leader-follower – This system of stocking management goes by several names. It is a method of utilizing two or more groups of animals, usually with different nutritional requirements, to graze sequentially on the same land area. The animal group with the highest nutritional need would have first access to a paddock. The groups that follow would have lower nutritional requirements.

Example: You just weaned your lambs. Now there are two groups of animals to manage; the dry ewes and the growing lambs. The lambs have the higher nutritional requirement and would benefit from being allowed first access to a paddock. The ewes with lower nutritional requirements could follow and would eat the lower quality forage not consumed by the lambs.




Direction of animal movement

Glossary




Life cycle – In the normal life cycle, worm eggs passed in the feces (manure) develop into larvae that are ingested by the sheep, and the larvae develop into adult worms in the sheep's abomasum or intestines. Egg hatching and larval development are dependent on warm temperatures and adequate moisture. The entire life cycle of *Haemonchus contortus*, perhaps the most important worm for us to manage, takes approximately 23-24 days under optimum conditions. The time needed for eggs to reach the infective larva stage can be as quick as 4 days but often is about 7-10 days. We can use this to our advantage in grazing strategies such as strip grazing with a back fence by moving the animals to a fresh strip of forage every 3-4 days. This means the animals move away from infective larvae developing on the pasture.



Glossary



Non-persistent dewormer – Most of the FDA-approved dewormers for sheep and goats are eliminated from the body relatively quickly, and their activity against adult and larval stages of worms is limited to a few hours or a day or so. In contrast, moxidectin, the active ingredient of Cydectin® drench, has a prolonged period of activity against worm larvae that are ingested with pasture forages (perhaps as long as 35 days if resistance has not developed to this chemical on your farm). From the sheep or goat's standpoint, this is a good thing as it may provide a prolonged period of protection against new worm infections. However, because the concentration of this chemical gradually declines over several weeks to less than lethal levels, research has shown that this may select for worms with resistance to this chemical. This has been shown with other dewormers with prolonged or persistent activity in other countries. This means that moxidectin should be used carefully in grazing animals to reduce selection pressure for resistance.



This graph shows the relative time for ivermectin and moxidectin, given orally, to be eliminated from sheep tissues.

Glossary




Refugia – the proportion of a farm’s parasitic worm population that escapes exposure to dewormers when animals are treated and that contributes to future worm generations on that farm. An example of parasite refugia is the worm larvae stages existing on the pasture that have developed from eggs. These larvae can survive through much of the grazing season, and even through typical winter weather, on the pastures. They are the life form of parasites that develop into a new generation of worms after they are consumed. Another example is the adult worms in untreated animals that are using the same pastures as the treated animals. Resistance to dewormers in the worm is a genetic trait that is passed on to future worm generations. Treatment with a dewormer removes the susceptible worms from an animal leaving those with genes for resistance to the dewormer to pass on that trait to future generations. The worms “in refugia” are available to mate with worms which survive treatment and thus help dilute the resistance genes. This is perhaps the most important concept in modern parasite control strategies to understand if we want sheep and goat farming to remain sustainable in the face of increasing resistance of worms to the dewormers we now have as well as any new ones we may get in the future.

Glossary



Rotational grazing – (or rotational stocking) is a method that utilizes recurring periods of grazing and rest among three or more paddocks in a grazing management unit throughout the time when grazing is allowed (Allen, et. al., 2011). Animals are restricted to a smaller pasture (paddock) for a limited time then removed and sent to a different pasture. Usually the movement is based on the amount of forage available in the fields. Typically animals may graze a field for three to seven days and then rotate to another field. Fields can be revisited when the forage re-grows enough to provide the appropriate amount of feed. The type of forage, the target amount of forage desired, how much residual leaf remained when the animals left and the weather while the paddock recovered all impact the time it takes for the paddock to be ready to re-graze. The paddock may be ready to re-graze within 14 to 45 days.



Glossary



Safe pasture – A safe pasture is one on which infective worm larvae are not present or are present in very low numbers. Pastures where sheep or goats have not grazed for the past year are usually safe because worm larvae can only exist for a defined period of time on their stored energy. Conventional tillage of the soil for row crop farming or planting annual forages effectively destroys worm larvae. Hayfields grazed in summer or fall after one or two harvests should have very low numbers of worm larvae even if the hayfield was grazed the previous fall. And pastures previously grazed by another species, such as horses or cattle, will be safe for sheep and goats because they do not share the same worms. Sheep, goats, and llamas do share the same worm species.



These goats and sheep are grazing on turnips which can provide good nutrition as well as a larvae-free place to graze.

Glossary



Set stocking – (sometimes called continuous grazing) is a method that allows a specific, non-variable number of animals on a specific, non-variable area of land during the time when grazing is allowed. For example if you had a pasture and turned all your animals into it at the start of spring and left them there for the whole grazing season, you would be set stocking.

A set number of animals have unrestricted access to this field only for the whole grazing season.

Glossary



Stocking density – is the number of pounds of live animals per acre of pasture or forage at one point in time. That takes the class and the size of animals out of the description. This is different from “stocking rate” which is the number of animals per acre over a time period (for example – 5 ewes per acre per year).

Example: You have 20 ewes weighing 150 lb. each for a total weight of 3,000 lb. You give them access to a one acre pasture. Your stocking density is 3,000 lb/ac. If you split the field in half and put all the animals on it, your stocking density is 6,000 lb/ac. If you gave them only one tenth of an acre, the stocking density would be 30,000 lb/ac. (1/10 acre equals a 66' x 66' square). If you want to get to 300,000 pounds of live animal per acre, then you must offer them one hundredth of an acre or a 21'x 21' area. This very high density is often referred to as “ultra-high stock density stocking” or “mob grazing” .

As you increase stocking density the amount of forage available for grazing becomes more critical. In addition, pasture contamination with worm larvae can become more serious. More frequent moves will be required to keep the livestock from over grazing the pasture, but several months of time may be required before worm larvae on the pasture disappear.

Manure distribution becomes more uniform as the stocking density increases. A more uniform distribution of manure also means a more uniform distribution of parasite eggs.



Glossary



Stocking density – is the number of pounds of live animals per acre of pasture or forage at one point in time. That takes the class and the size of animals out of the description. This is different from “stocking rate” which is the number of animals per acre over a time period (for example – 5 ewes per acre per year).




These goats are grazing a large acreage of southern Ohio hillside. Manure deposition, and worm larvae concentration, will be reduced by this low stocking density, and the risk of parasitism is much reduced.

Glossary



Strip grazing – is defined as a method that confines animals to an area of grazing land to be grazed in a relatively short time. The strip size is varied to allow access to a specific land area. Typically this method is used when more efficient utilization of the standing forage is desired. It limits access to forage with temporary fencing. If the forage has the potential to re-grow, a back fence would be used to prevent overgrazing the portion already grazed. The size of paddock could change across the field if the amount of forage is not uniform in the whole field.



To continue navigation, select "Next" button





Strip grazing



Sheep have just finished grazing the forage in the strip on the left and will be moved to fresh forage at the right of the picture. In this case they are grazing standing corn.

Glossary



Summer annual – plants that are planted in the spring for grazing during that same growing season. They sprout, flower, produce seed, and die during the warmer months of the year. Examples include sorghum/sudangrass hybrids, pearl millet, forage soybeans, cowpeas, annual sericea lespedeza, and even corn. Turnips and rape, plants in the Brassica family, are not true summer annual plants but are often planted and used as summer annuals in the northern region of the USA.




These animals are grazing brown mid-rib sorghum/sudangrass, a rapidly growing summer annual plant. Infective worm larvae are unlikely to be found on the leaves of this plant as they are too high off the ground. However, when the sorghum/sudangrass fields are re-grazed, they may be found on annual grasses and other plants growing along with this plant if the animals previously grazing the plants were infected and shedding worm eggs.

Glossary



TST – “Targeted Selective Treatment” – Treatment of only those animals that will most benefit from treatment, leaving the rest of the flock or herd untreated. Worm numbers in sheep and goats are not uniform across members of the flock or herd with approximately 70-80% of the worms found in only 20–30% of the animals; the majority of the animals have relatively low worm burdens. TST strategies are directed toward the animals that are clinically affected by parasites, those animals most susceptible to disease, or those that are likely to contaminate the pasture the most. Examples of criteria that can be used to selectively choose animals for treatment include anemia (the FAMACHA© system); thin body condition; reduced live weight gains; elevated worm egg counts in feces; and below average milk production.



This graph represents the fecal worm egg counts in a group of lambs experiencing severe clinical parasitism from *Haemonchus contortus*. Just 10 (21%) of these 46 lambs were excreting about 77% of all the eggs in the group. If they can be identified, such as with FAMACHA©, the animals most needing treatment and which contribute most to pasture contamination can be treated. The remaining animals contribute little to pasture worm contamination and that contamination will be from unselected worms – thus leaving a refugia to dilute the genes from resistant worms.

Glossary



Winter annual – In the context of a grazing system, these are plants that are planted in the fall and intended for grazing the following spring. They live one year. Examples include wheat and cereal rye.



These animals are grazing cereal rye. A winter annual such as this can provide larvae-free pasture in the spring.

Additional Resources



PUBLICATIONS:

Allen V.G., C. Batello, E.J. Berretta, J. Hodgson, M. Kothmann, X. Li, J. McIvor, J. Milne, C. Morris, A. Peeters , & M. Sanderson (2011). [An international terminology for grazing lands and grazing animals](#). *Grass and Forage Science*, 66, 2–28.

Coffey L. [Tools for Managing Internal Parasites in Small Ruminants: Animal Selection](#). National Center for Appropriate Technology, Publication IP400, 2012.

Coffey L., Hale M., Terrill T., Mosjidis J., Miller J., & Burke J. [Tools for Managing Internal Parasites in Small Ruminants: Sericea Lespedeza](#). National Center for Appropriate Technology, Publication IP316, 2007.

Hale M., Burke J., Miller J., & Terrill T. [Tools for Managing Internal Parasites in Small Ruminants: Copper Wire Particles](#). National Center for Appropriate Technology, Publication IP317, 2007.

LINKS:

[American Consortium for Small Ruminant Parasite Control](#)

[OSU Sheep Team](#)

[OSU Veterinary Extension](#)

[Veterinary Preventive Medicine Fact Sheet Index](#)



Return to [Table of Contents](#) 28

Determining Dewormer Effectiveness



Because some reliance on chemical dewormers is necessary for most farms, it is very important to know how well these products perform against the worms on your farm.

Do you have data indicating that at least one dewormer remains **highly effective** on your farm? (“data” means fecal egg count reduction testing or DrenchRite® Assay results)

yes

no



Determining Dewormer Effectiveness



I have data showing at least one highly effective dewormer available for my farm:

Stay with the dewormer that works, use it with a system that allows maintenance of a refugia of unselected worms on your farm, and continue to monitor effectiveness at least every 2 years.

It is a good idea to monitor for deworming success occasionally between formal testing for efficacy. This can be done by collecting manure samples from animals that were dewormed 10-14 days previously and having your veterinarian determine the worm egg count. Preferably, 15 samples should be collected and a quantitative method to count eggs used.



Manure, or feces, samples can be easily collected by inverting a resealable plastic storage bag over the hand and using a finger to gently pull some manure into the bag. It can then be turned right side out and the animal's identification placed on it with a permanent marker.

See: [fecal egg count reduction testing](#)



Return to [Table of Contents](#) 30

Determining Dewormer Effectiveness



No data showing highly effective dewormers on my farm:

Check for dewormer resistance on your farm.


- [Fecal egg count reduction testing](#):

- Consult your veterinarian for fecal egg count reduction testing.

- [DrenchRite® Assay](#):

- Consult the [laboratory](#) at the University of Georgia for DrenchRite® Assay.

Now that you know the resistance status of your flock, please select the “Next” button. 



[McMaster slide](#)



Determining Dewormer Effectiveness



Does one or more FDA-approved dewormer exist for your animal species that should be effective?

yes

no

FDA-approved dewormers available for use in sheep and goats in the US as of August, 2012

| Drug class | Active ingredient | Trade names |
|--|-------------------|--|
| Benzimidazole | Albendazole | Valbazen ^{®1} |
| | Fenbendazole | Safeguard ^{®2} |
| Imidazothiazole and Tetrahydropyrimidine | Levamisole | Prohibit ^{®1} , Levasol ¹ , Tramisol ^{®1} |
| | Morantel tartrate | Rumatel ^{®3} |
| Macrocyclic lactones ivermectin | Ivermectin | Ivomec Drench for Sheep ^{®1} |
| | Moxidectin | Cydectin ^{®1} |

¹FDA-approved for use in sheep (not for sheep producing milk for human consumption)

²FDA-approved for use in goats (not for goats producing milk for human consumption)

³FDA-approved for use in goats (feed supplement only; is approved for lactating does)

Other dewormers are FDA-approved for other species and might be used in sheep and goats, and some products in the table above can be used in both sheep and goats. However, all such uses are considered extra-label and must be by veterinary prescription.

Determining Dewormer Effectiveness



No effective FDA-approved dewormers:

Consult your veterinarian to consider the use of dewormers approved for other species or possibly copper oxide wires.



Determining Dewormer Effectiveness



Yes-one or more effective FDA-approved dewormers exist:

Is Cydectin® (moxidectin) one of the dewormers that should be effective?

yes

no



Determining Dewormer Effectiveness



Cydectin[®] is an effective FDA-approved dewormers:

Is Cydectin[®] (moxidectin) the only dewormer that is effective?

yes

no




Determining Dewormer Effectiveness



Cydectin® is the only effective FDA-approved dewormer:

Reserve moxidectin use to selective deworming strategies (such as the [FAMACHA©](#) system) or strategies that allow a "[refugia](#)" of unselected parasites in order to prolong its useful life.



Determining Dewormer Effectiveness



Cydectin® is not the only effective FDA-approved dewormer:

Reserve moxidectin use until others fail or reserve its use to feedlot animals that will not go to pasture after deworming.



Lactating Females



Lactating females with nursing lambs and kids.

Will these animals be grazing?

yes

no



Lactating Females



No grazing:

Lactating females kept indoors or on dry lot will get no exposure to [GIN](#).



Lactating Females



Yes-grazing season:

Season being grazed?

Spring

Summer

Fall

Winter

Lactating Females



Spring:

What type of forage is being grazed?

Permanent pasture

**Winter annual
(cereal rye, wheat)**

Hayfield

Lactating Females



Spring-permanent pasture:

When was the permanent pasture last grazed by sheep or goats?

Previous Spring

Previous Summer

Previous Fall

Previous Winter

Same Spring

Lactating Females



Spring-permanent pasture-previous spring/previous summer:

Pastures grazed in the spring that were last grazed the previous summer or spring should be relatively worm larvae-free. Worm larvae exposed to summer weather will use up their stored energy and be unlikely to survive winter.



Lactating Females



Spring-permanent pasture-previous fall:

Ewes and lambs grazing these pastures will likely be exposed to overwintered parasite larvae. Use [FAMACHA©](#) every two weeks or more frequently if your options for grazing are limited. Consult OSUE Fact Sheet VME-28-12 for more information on managing springtime pastures with lactating females and lambs and kids and especially for larger flocks.

See OSUE Fact Sheet VME-28-12 [“Strategies for Coping with Parasite Larvae on Pastures in the Springtime in Ohio”](#)



Lactating Females



Spring-permanent pasture-previous winter:

Worm eggs that may have been deposited on the pasture during winters where freezing temperatures are common do not survive well. The first two stages of larvae developing from eggs that do hatch during periods of warm weather in winter are very susceptible to drying. Third stage larvae are the infectious stage and survive winter much better. The risk of pastures grazed during winter will depend on average temperatures in your area, moisture, number of animals grazing, and their egg output.



Lactating Females



Spring-permanent pasture-same spring:

In addition to exposure to overwintered larvae, ewes and lambs will be exposed to larvae developing from eggs shed in an earlier grazing this spring. If some strategy to minimize egg shedding during the earlier grazing is not employed, these pastures can have high levels of worm larvae. Deworming only thin ewes and ewes with twins and triplets can reduce pasture contamination while still providing a [refugia](#) of worms not exposed to dewormer.

See OSUE Fact Sheet VME-28-12 [“Strategies for Coping with Parasite Larvae on Pastures in the Springtime in Ohio”](#)



Lactating Females



Spring-winter annual:

Winter annuals are usually planted after tillage of the soil and are usually planted in the fall following harvest of a crop. If they were planted on fields where sheep or goats had grazed that season, tillage will destroy most worm larvae making these pastures safe. If a “no-till” cultivation was used to plant the winter annual, some larvae may survive. No data exists to predict how many, but it is likely to be only small numbers.



Lactating Females



Spring-hayfield:

If hayfields can be grazed in the spring, thus effectively harvesting “the first cutting” with the sheep, they usually provide a worm larvae-free place to put the animals. If combined with [strip grazing](#) and back fencing, hayfield grazing can be even more useful for worm control and be more efficient in forage usage. It takes at least 3-4 days under the most ideal weather conditions for a worm egg to hatch and reach the infective larva stage. If the fences are moved across the clean hayfield at 3-4 day intervals, and the back fence prevents sheep from grazing where eggs may have been deposited, both ewes and growing lambs will not acquire new infections as long as these larvae-free fields are available. Given the difficulty of making good quality first cutting hay in Ohio because of typical weather conditions, this can be a very efficient way of utilizing the forage. If the hay is predominantly alfalfa or clover, a strategy to manage bloat will need to be developed. If the hayfield is not grazed again during the summer or fall, it should be larvae free by the next spring because most infective larvae will use up their stored energy over the summer, and hay making will expose them to drying out. If strip grazing is not possible, deworming the lambs before moving to the clean field, leaving all or most of the ewes untreated, will provide the lambs with some protection against infection. In such situations, infective worm larvae will begin to accumulate to significant levels on the pastures by 30-40 days of grazing. This may not be an important issue on hayfields in which one expects later harvest as it is unlikely one would wish to graze it that long.



Lactating Females



Summer:

What type of forage is being grazed?

Sericea lespedeza

Summer annual

Permanent pasture

Lactating Females



Summer-Sericea lespedeza:

Information about planting and using sericea lespedeza can be found at the website for the [American Consortium for Small Ruminant Parasite Control](#).



Lactating Females



Summer-summer annual:

Summer annuals such as turnips, kale, or sorghum/sudangrass hybrids can provide a worm larvae-free place to graze ewes and preweaned lambs. These are most effectively grazed using strip grazing with a back fence to prevent animals having access to previously grazed portions of the field. If the fences are moved across these clean pastures at 3-4 day intervals, and the back fence prevents sheep from grazing where eggs may have been deposited, both ewes and growing lambs will not acquire new infections as long as these larvae-free fields are available. If strip grazing is not possible, deworming the lambs before moving to the clean field, leaving all or most of the ewes untreated, will provide the lambs with some protection against infection and still provide a refugia of unselected worms to help reduce selection for dewormer resistance.

See OSUE Fact Sheet VME-28-12 ["Strategies for Coping with Parasite Larvae on Pastures in the Springtime in Ohio"](#)



Lactating Females



Summer-permanent pasture:

Permanent pastures for lactating ewes and lambs can be very dangerous if they were grazed in the spring with lactating ewes shedding large numbers of parasite eggs. Consider [TST](#) of ewes in spring and [FAMACHA©](#) every 7-14 days throughout the summer. Also consider alternate species grazing, annual forages, or hayfield grazing to create larvae-free places to graze.

See OSUE Fact Sheet VME-28-12 [“Strategies for Coping with Parasite Larvae on Pastures in the Springtime in Ohio”](#)



More information about the FAMACHA© system can be found at: <http://www.sheepandgoat.com/ACSRP/C/Resources/famacha.html>

FAMACHA© cards and training may be available from your veterinarian or call your Extension educator to request training in your area.

Lactating Females



Fall:

What type of forage is being grazed?

Permanent Pasture

Hayfield

Lactating Females



Fall-permanent pasture:

Permanent pastures will likely have significant worm larvae on them if grazed during the spring and summer depending on the class of animals grazing them. Fall lambing ewes, or ewes nursing late spring lambs, grazing these pastures will need to be monitored, preferably with [FAMACHA®](#), until the weather gets cold enough to retard transmission (ground temps below 50°F). Lambs older than about 6 weeks-of-age should also be monitored.



Lactating Females



Fall-hayfield:

Hayfields not previously grazed during this grazing season can provide a worm larvae-free place to graze sheep in the fall. [Strip grazing](#) with backfencing is the preferred technique. If alfalfa or clover is used, a bloat prevention strategy will need to be used. Care must be taken to allow the forage to recover before winter if the field is to be used for hay the following year.



**strip grazing with a back fence –
minimum 3.5 days to go from
worm egg to infective L₃ on
forage**

Lactating Females



Winter:

Good options in some areas include standing corn, harvested corn fields or other crop aftermath, and stockpiled fescue. Grass pastures previously grazed in the summer and fall, such as stockpiled fescue, may contain worm larvae. Transmission can occur if the animals are forced to eat low to the ground or if the weather is warm for extended periods. Nutritional needs of the ewes and lambs must be met. Supplemental feeding of ewes may be necessary on some forages.



Weaned Growing Lambs and Kids



Weaned, growing lambs and kids.

Will these animals be grazing?

yes

no

Weaned Growing Lambs and Kids



No grazing:

Weaned lambs and kids kept indoors or on dry lot will get no exposure to [GIN](#). If they have not been on pastures where other sheep have grazed, they do not need dewormed.

The important GIN in sheep and goats do not easily complete their [life cycle](#) in dry lot or barn environments.



Weaned Growing Lambs and Kids



Yes-grazing season:

Season being grazed?

Spring*

Summer

Fall

Winter

** It is unusual to graze weaned lambs in early spring on most farms in the North Central region of the US.*

Return to [Table of Contents](#) 59

Weaned Growing Lambs and Kids



Spring:

What type of forage is being grazed?

Permanent pasture

**Winter annual
(cereal rye, wheat)**

Hayfield

Weaned growing lambs and kids.



Spring-permanent pasture:

When was the permanent pasture last grazed by sheep or goats?

Previous Spring

Previous Summer

Previous Fall

Previous Winter

Same Spring

Weaned growing lambs and kids.



Spring-permanent pasture-previous spring/previous summer:

In temperate regions, pastures grazed in the spring that were last grazed the previous summer or spring should be relatively worm larvae-free. Worm larvae exposed to summer weather will use up their stored energy and be unlikely to survive winter. In warmer climates of the southeastern USA, there will be fewer larvae but some infective larvae may be found year-round.



Weaned Growing Lambs and Kids



Spring-permanent pasture-previous fall:

Weaned lambs grazing these pastures will likely be exposed to overwintered parasite larvae. Use the [FAMACHA©](#) system every two weeks or more frequently if your options for grazing are limited. Consult OSUE Fact Sheet VME-28-12 for more information on managing springtime pastures with lactating ewes and lambs and especially for larger flocks.

See OSUE Fact Sheet VME-28-12 [“Strategies for Coping with Parasite Larvae on Pastures in the Springtime in Ohio”](#)



Weaned Growing Lambs and Kids



Spring-permanent pasture-previous winter:

Worm eggs that may have been deposited on the pasture during winters where freezing temperatures are common do not survive well. The first two stages of larvae developing from eggs that do hatch during periods of warm weather in winter are very susceptible to drying. Third stage larvae are the infectious stage and survive much better. The risk of pastures grazed during winter will depend on average temperatures in your area, moisture, number of animals grazing, and their egg output.



Weaned Growing Lambs and Kids



Spring-permanent pasture-same spring:

In addition to exposure to overwintered larvae, these animals will be exposed to larvae developing from eggs shed in an earlier grazing this spring. If some strategy to minimize egg shedding during the earlier grazing was not employed, these pastures can have high levels of worm larvae. Deworming only thin ewes and ewes with twins and triplets can reduce pasture contamination while still providing a [refugia](#) of worms not exposed to dewormer.

See OSUE Fact Sheet VME-28-12 [“Strategies for Coping with Parasite Larvae on Pastures in the Springtime in Ohio”](#)

Lactating ewes and does can excrete large numbers of worm eggs. The larvae developing from them are a major source of infection for the lambs.



Weaned Growing Lambs and Kids



Spring-winter annual:

Winter annuals are usually planted after tillage of the soil and are usually planted in the fall following harvest of a crop. If they were planted on fields where sheep or goats had grazed that season, tillage will destroy most worm larvae making these pastures safe. If a “no-till” cultivation was used to plant the winter annual, some larvae may survive. No data exists to predict how many but it is likely to be only small numbers.



Weaned Growing Lambs and Kids



Spring-hayfield:

If hayfields can be grazed in the spring, thus effectively harvesting “the first cutting” with the sheep, they usually provide a worm larvae-free place to put the animals. If combined with [strip grazing](#) and back fencing, hayfield grazing can be even more useful for worm control and be more efficient in forage usage. It takes at least 3-4 days under the most ideal weather conditions for a worm egg to hatch and reach the infective larva stage. If the fences are moved across the clean hayfield at 3-4 day intervals, and the back fence prevents the animals from grazing where eggs may have been deposited, lambs will not acquire new infections as long as these larvae-free fields are available. Given the difficulty of making good quality first cutting hay in our region because of typical weather conditions, this can be a very efficient way of utilizing the forage. If the hay is predominantly alfalfa or clover, a strategy to manage bloat will need to be developed. If the hayfield is not grazed again during the summer or fall, it should be larvae-free by the next spring because most infective larvae will use up their stored energy over the summer, and hay making will expose them to drying out. If the lambs and kids were born in a barn in winter and placed on hayfields that were not grazed the previous fall, they should stay worm free as long as they are on the hayfield. Lambs and kids do not acquire the important [GIN](#) in a typical barn environment and will not acquire worms on pastures that have not been previously grazed.

Weaned Growing Lambs and Kids



Summer:

What type of forage is being grazed?

Sericea lespedeza

Summer annual

Permanent pasture

Weaned Growing Lambs and Kids



Summer-sericea lespedeza:

Information about planting and using sericea lespedeza can be found at the website for the [American Consortium for Small Ruminant Parasite Control](#).



Weaned Growing Lambs and Kids



Summer-summer annual:

[Summer annuals](#) such as turnips, kale, millet, cowpeas, or sorghum/sudangrass hybrids can provide a worm larvae-free place to graze growing lambs. These forages are most effectively grazed using [strip grazing](#) with a back fence to prevent animals having access to previously grazed portions of the field. If the fences are moved across these clean pastures at 3-4 day intervals, and the back fence prevents the animals from grazing where eggs may have been deposited, growing lambs and kids will not acquire NEW infections as long as these larvae-free fields are available. If strip grazing is not possible, deworming the animals before moving to the clean field will provide them with some protection against infection. Avoid treating all the animals and immediately moving them to a worm larvae-free field as this will result in selection for dewormer resistant worms.

This selection can be reduced by: 1) deworming the group but leaving the heaviest 10-25% of lambs untreated; 2) treating all the animals with a [non-persistent dewormer](#) while still on contaminated pasture and waiting 5-7 days before moving them to allow them to acquire a small infection with unselected worms; or 3) moving to the clean pasture and waiting 5-7 days before treating them. It is important when using summer annuals for growing lambs and kids to make sure their nutritional needs are met.

See OSUE Fact Sheet VME-28-12 [“Strategies for Coping with Parasite Larvae on Pastures in the Springtime in Ohio”](#)



Return to [“Season being grazed”](#) 70

Weaned Growing Lambs and Kids



Summer-permanent pasture:

Permanent pastures grazed in summer with weaned lambs and kids can be very dangerous if they were grazed in the spring with lactating females shedding large numbers of parasite eggs. Consider [TST](#) of lactating females in spring to reduce egg shedding. If only permanent pastures previously grazed by sheep or goats are available, consider using the [FAMACHA©](#) scoring system every 7-14 days throughout the summer for weaned lambs and kids. If other options might exist, consider alternate species grazing, grazing summer annual forages, or hayfield grazing to create larvae-free places to graze for lambs.

See OSUE Fact Sheet VME-28-12 [*"Strategies for Coping with Parasite Larvae on Pastures in the Springtime in Ohio"*](#)

More information about the FAMACHA© system can be found at:
<http://www.sheepandgoat.com/ACSRPC/Resources/famacha.html>

FAMACHA© cards and training may be available from your veterinarian or call your Extension educator to request training in your area.

Weaned Growing Lambs and Kids



Fall:

What type of forage is being grazed?

Permanent Pasture

Hayfield

Weaned Growing Lambs and Kids



Fall-permanent pasture:

Permanent pastures will likely have significant numbers of worm larvae on them if grazed during the spring and summer depending on the class of animals grazing them. Growing lambs and kids grazing these pastures will need to be monitored, preferably with [FAMACHA®](#), until the weather gets cold enough to retard transmission (ground temps below 50°F). Other forms of [TST](#) may also be appropriate especially in larger flocks and herds.



Weaned Growing Lambs and Kids



Fall-hayfield:

Hayfields not previously grazed this grazing season can provide a worm larvae-free place to graze lambs in the fall. [Strip grazing](#) with backfencing is the preferred technique. If alfalfa or clover is used, a bloat prevention strategy will need to be used. Care must be taken to allow the forage to recover before winter if the field is to be used for hay the following year.



Lambs grazing alfalfa in mid August.

Weaned Growing Lambs and Kids



Winter:

Good options in some areas include standing corn, harvested corn fields, and stockpiled fescue. Grass pastures grazed in the summer and fall, such as stockpiled fescue, may contain worm larvae. Transmission can occur if the animals are forced to eat low to the ground or if the weather is warm for extended periods. Nutritional needs of the animals must be met if good gains are expected. In warmer regions, winter annuals (wheat, rye, vetch) can be grazed.



Replacement Females



Replacement females.

Will these animals be grazing?

yes

no

Replacement Females



No grazing:

Replacement females kept indoors or on dry lot will get no exposure to [GIN](#).



Replacement Females



Yes grazing:

Will they be grazed with lactating or dry ewes?

yes

no

Replacement Females



Yes-grazing with lactating or dry ewes/does:

Manage parasitism for replacement females the same as lactating or dry ewes/does that they graze with.

- Go to [Lactating Females](#)
- Go to [Dry Ewes and Does](#)



Replacement Females



Yes grazing-not grazing with lactating or dry ewes/does-grazing season:

Season being grazed?

Spring

Summer

Fall

Winter

Replacement Females



Spring:

What type of forage is being grazed?

Permanent pasture

**Winter annual
(cereal rye, wheat)**

Hayfield

Replacement Females



Spring-permanent pasture:

When was the permanent pasture last grazed by sheep or goats?

Previous Spring

Previous Summer

Previous Fall

Previous Winter

Same Spring

Replacement Females



Spring-permanent pasture-previous spring/previous summer:

Pastures grazed in the spring that were last grazed the previous summer or spring should be relatively worm larvae-free. Worm larvae exposed to summer weather will use up their stored energy and be unlikely to survive winter.

Replacement Females



Spring-permanent pasture-previous fall:

Although these pastures will likely have overwintered parasite larvae, replacement females that have had one season of grazing will have acquired some immunity to parasitism, and they can withstand a worm challenge more effectively than lactating ewes and growing lambs. If they are in good body condition and pasture nutritive value is good, risk of severe internal parasitism is low. It would be a good idea to monitor body condition and eyelid score occasionally ([FAMACHA©](#) system). Consult OSUE Fact Sheet VME-28-12 for more information on managing springtime pastures for replacement females.

See OSUE Fact Sheet VME-28-12 [*“Strategies for Coping with Parasite Larvae on Pastures in the Springtime in Ohio”*](#)

FAMACHA© cards and training may be available from your veterinarian or call your Extension educator to request training in your area.



Replacement Females



Spring-permanent pasture-previous winter:

Worm eggs that may have been deposited on the pasture during winters where freezing temperatures are common do not survive well. The first two stages of larvae developing from eggs that do hatch during periods of warm weather in winter are very susceptible to drying. Third stage larvae are the infectious stage and survive winter much better. The risk of pastures grazed during winter will depend on average temperatures in your area, moisture, number of animals grazing, and their egg output.



Replacement Females



Spring-permanent pasture-same spring:

In addition to exposure to overwintered larvae, these animals will be exposed to larvae developing from eggs shed in an earlier grazing this spring. If some strategy to minimize egg shedding during the earlier grazing is not employed, these pastures can have high levels of worm larvae. Deworming only thin ewes and ewes with twins and triplets can reduce pasture contamination while still providing a [refugia](#) of worms not exposed to dewormer. Non-lactating replacement females will be more resistant to clinical parasitism than lactating females or growing lambs.

Replacement Females



Spring-winter annual:

Winter annuals are usually planted after tillage of the soil and are usually planted in the fall following harvest of a crop. If they were planted on fields where sheep or goats had grazed that season, tillage will destroy most worm larvae making these pastures safe. If a “no-till” cultivation was used to plant the winter annual, some larvae may survive. No data exists to predict how many but it is likely to be only small numbers.



Replacement Females



Spring-hayfield:

If hayfields can be grazed in the spring, thus effectively harvesting “the first cutting” with the sheep, they usually provide a worm larvae-free place to put the animals. If combined with [strip grazing](#) and back fencing, hayfield grazing can be even more useful for worm control and be more efficient in forage usage. It takes at least 3-4 days under the most ideal weather conditions for a worm egg to hatch and reach the infective larva stage. If the fences are moved across the clean hayfield at 3-4 day intervals, and the back fence prevents the animals from grazing where eggs may have been deposited, animals will not acquire new infections as long as these larvae-free fields are available. Given the difficulty of making good quality first cutting hay in our region because of typical weather conditions, this can be a very efficient way of utilizing the forage. If the hay is predominantly alfalfa or clover, a strategy to manage bloat will need to be developed. If the hayfield is not grazed again during the summer or fall, it should be larvae-free by the next spring because most infective larvae will use up their stored energy over the summer, and hay making will expose them to drying out.

Replacement Females



Summer:

What type of forage is being grazed?

Sericea lespedeza

Summer annual

Permanent pasture

Replacement Females



Summer-sericea lespedeza:

Information about planting and using sericea lespedeza can be found at the website for the [American Consortium for Small Ruminant Parasite Control](#).



Replacement Females



Summer-summer annual:

Summer annuals such as turnips, kale, millet, cowpeas, or sorghum/sudangrass hybrids can provide a worm larvae-free place to graze replacement females although they are better able to withstand a parasite challenge than young growing lambs. These forages are most effectively grazed using **strip grazing** with a back fence to prevent animals having access to previously grazed portions of the field. If the fences are moved across these clean pastures at 3-4 day intervals, and the back fence prevents sheep from grazing where eggs may have been deposited, animals will not acquire NEW infections as long as these larvae-free fields are available. If strip grazing is not possible and you will be grazing replacement females that you suspect have a worm burden sufficient to retard their growth, deworming the animals before moving to the clean field will provide the them with protection against an increasingly serious infection. Avoid treating all the animals and immediately moving them to a worm larvae-free field as this will result in selection for dewormer resistant worms. This selection can be reduced by 1) deworming the group but leaving the heaviest 10-25% of animals untreated; 2) treating all the animals with a non-persistent dewormer while still on contaminated pasture and waiting 5-7 days before moving them to allow them to acquire a small infection with unselected worms; or 3) moving to the clean pasture and waiting 5-7 days before treating them. It is important when using summer annuals for growing animals to make sure their nutritional needs are met.

Replacement Females



Summer-permanent pasture:

Permanent pastures grazed in summer can have very high levels of parasite larvae on them if they were grazed in the spring with lactating ewes or growing lambs that were shedding large numbers of parasite eggs. Although non-lactating replacement females should have a higher level of ability to withstand a parasite challenge than lambs in their first grazing season, they should be monitored for signs of parasitism. If only permanent pastures previously grazed by sheep are available, consider using the [FAMACHA©](#) scoring system every 14 days throughout the summer to monitor. Alternatively, monitoring body condition and group level [fecal egg counts](#) may provide an alternative approach. If treatment is needed consider using the FAMACHA© system as a guide for which animals to be treated or consider using a [TST](#) approach where the group is dewormed but the heaviest 25% of animals remain untreated. If other options might exist, consider alternate species grazing, grazing summer annual forages, or hayfield grazing to create larvae-free, or reduced risk, places to graze because you will want these animals to reach breeding condition without setback.

Replacement Females



Fall:

What type of forage is being grazed?

Permanent Pasture

Hayfield

Replacement Females



Fall-permanent pasture:

Permanent pastures may have significant numbers of worm larvae on them if they were grazed during the spring and summer by lactating ewes or growing lambs and depending on the success of parasite control strategies. Replacement females grazing these pastures should be monitored, with [FAMACHA®](#), body condition scoring, or [fecal egg counts](#), until the weather gets cold enough to retard transmission (ground temps below 50°F).

Lactating ewes and does can excrete large numbers of worm eggs. The larvae developing from them are a major source infection for the lambs and provide contamination for other sheep or goats grazing the pastures.



Replacement Females



Fall-hayfield:

Hayfields not previously grazed this grazing season can provide a worm larvae-free place to graze in the fall and may provide excellent nutrition for flushing ewes. [Strip grazing](#) with backfencing is the preferred technique. If alfalfa or clover is used, a bloat prevention strategy will need to be used. Care must be taken to allow the forage to recover before winter if the field is to be used for hay the following year.



**strip grazing with a back fence –
minimum 3.5 days to go from
worm egg to infective L₃ on
forage**

Replacement Females



Winter:

Replacement females grazing in winter will be unlikely to acquire many worms if it is cold. Good options in some areas include standing corn, harvested corn fields, and stockpiled fescue.



Dry Ewes and Does



Dry ewes and does.

Will these animals be grazing?

yes

no

Dry Ewes and Does



No grazing:

Dry ewes kept indoors or on dry lot will get no exposure to [GIN](#).



Dry Ewes and Does



Yes-grazing season:

Season being grazed?

Spring

Summer

Fall

Winter

Dry Ewes and Does



Spring:

What type of forage is being grazed?

Permanent pasture

**Winter annual
(cereal rye, wheat)**

Hayfield

Dry Ewes and Does



Spring-permanent pasture:

When was the permanent pasture last grazed by sheep or goats?

Previous Spring

Previous Summer

Previous Fall

Previous Winter

Same Spring

Dry Ewes and Does



Spring-permanent pasture-previous spring/previous summer:

Pastures grazed in the spring that were last grazed the previous summer or spring should be relatively worm larvae-free. Worm larvae exposed to summer weather will use up their stored energy and be unlikely to survive winter.

Dry Ewes and Does



Spring-permanent pasture-previous fall:

Although these pastures will likely have overwintered parasite larvae, non-lactating females that have had one season of grazing will have acquired some immunity to parasitism, and they can withstand a worm challenge more effectively than lactating ewes and growing lambs. If they are in good body condition and pasture nutritive value is good, risk of severe internal parasitism is low. It would be a good idea to monitor body condition and eyelid score occasionally ([FAMACHA©](#) system). Consult OSUE Fact Sheet VME-28-12 for more information on managing springtime pastures for dry ewes/does.

See OSUE Fact Sheet VME-28-12 [*“Strategies for Coping with Parasite Larvae on Pastures in the Springtime in Ohio”*](#)




Dry Ewes and Does



Spring-permanent pasture-previous winter:

Worm eggs that may have been deposited on the pasture during winters where freezing temperatures are common do not survive well. The first two stages of larvae developing from eggs that do hatch during periods of warm weather in winter are very susceptible to drying. Third stage larvae are the infectious stage and survive much better. The risk of pastures grazed during winter will depend on average temperatures in your area, moisture, number of animals grazing, and their egg output. However, non-lactating females can withstand some worm larval challenge without becoming severely parasitized. Use body condition score and/or the [FAMACHA©](#) system to monitor these animals.



STOP Return to "[Season being grazed](#)" 104

Dry Ewes and Does



Spring-permanent pasture-same spring:

These animals will be exposed to larvae developing from eggs shed in an earlier grazing this spring. If some strategy to minimize egg shedding during the earlier grazing was not employed, these pastures can have high levels of worm larvae. For example, if lactating females were the last to use this pasture, then deworming thin ewes and ewes with twins and triplets can reduce pasture contamination while still providing a [refugia](#) of worms not exposed to dewormer. However, non-lactating mature females will be more resistant to clinical parasitism than lactating females or growing lambs.

Non-lactating females seem to be most at risk for severe parasitism in the first few weeks after weaning. Watch thin animals closely or consider deworming animals in thin body condition after weaning. Ewes and does in good body condition at weaning probably will not require deworming.



Dry Ewes and Does



Spring-winter annual:

Winter annuals are usually planted after tillage of the soil and are usually planted in the fall following harvest of a crop. If they were planted on fields where sheep or goats had grazed that season, tillage will destroy most worm larvae making these pastures safe. If a “no-till” cultivation was used to plant the winter annual, some larvae may survive. No data exists to predict how many will survive, but it is likely to be only small numbers.



Dry Ewes and Does



Spring-hayfield:

If hayfields can be grazed in the spring, thus effectively harvesting “the first cutting” with the sheep, they usually provide a worm larvae-free place to put the animals. If combined with [strip grazing](#) and back fencing, hayfield grazing can be even more useful for worm control and be more efficient in forage usage. It takes at least 3-4 days under the most ideal weather conditions for a worm egg to hatch and reach the infective larva stage. If the fences are moved across the clean hayfield at 3-4 day intervals, and the back fence prevents the animals from grazing where eggs may have been deposited, animals will not acquire new infections as long as these larvae-free fields are available. Given the difficulty of making good quality first cutting hay in our region because of typical weather conditions, this can be a very efficient way of utilizing the forage. If the hay is predominantly alfalfa or clover, a strategy to manage bloat will need to be developed. If the hayfield is not grazed again during the summer or fall, it should be larvae-free by the next spring because most infective larvae will use up their stored energy over the summer, and hay making will expose them to drying out.

Dry Ewes and Does



Summer:

What type of forage is being grazed?

Sericea lespedeza

Summer annual

Permanent pasture

Dry Ewes and Does



Summer: sericea lespedeza:

Information about planting and using sericea lespedeza can be found at the website for the [American Consortium for Small Ruminant Parasite Control](#).



STOP Return to "[Season being grazed](#)" 109

Dry Ewes and Does



Summer-summer annual:

Summer annuals such as turnips, kale, millet, cowpeas, or sorghum/sudangrass hybrids can provide a worm larvae-free place to graze mature non-lactating females. However, they are much better able to withstand a parasite challenge than young growing lambs or lactating ewes. These forages are most effectively grazed using **strip grazing** with a back fence to prevent animals having access to previously grazed portions of the field. If the fences are moved across these clean pastures at 3-4 day intervals, and the back fence prevents sheep from grazing where eggs may have been deposited, animals will not acquire NEW infections as long as these larvae-free fields are available. Dry females may not need the high quality nutrition that some summer annuals provide. Sorghum Sudangrass hybrids may be ideally suited to non-lactating females. As with any other class of animals, avoid treating all the animals and immediately moving them to a worm larvae-free field as this will result in selection for dewormer resistant worms. Typically, after the lambs or kids are weaned, fecal worm egg counts fall to low levels unless the ewes/does have been stressed by poor quality nutrition and are in very thin body condition. They should be evaluated at weaning and treated selectively if needed. The forage quality should be sufficient for the ewes/does to regain body condition in preparation for breeding.

Dry Ewes and Does



Summer-permanent pasture:

Permanent pastures grazed in summer can have very high levels of parasite larvae on them if they were grazed in the spring with lactating females that were shedding large numbers of parasite eggs. Typically, after the lambs or kids are weaned, fecal worm egg counts fall to low levels unless the ewes/does have been stressed by poor quality nutrition and are in very thin body condition. They should be evaluated at weaning and treated selectively if needed. The forage quality should be sufficient for the ewes/does to regain body condition in preparation for breeding. Although mature non-lactating females are much better able to withstand a parasite challenge than young growing lambs, they should be monitored for signs of parasitism. If only permanent pastures previously grazed by sheep are available, consider using the [FAMACHA©](#) scoring system every 14 days throughout the summer to monitor. Alternatively, monitoring body condition and group level [fecal egg counts](#) may provide a different approach. If treatment is needed consider using the FAMACHA© system as a guide for which animals to be treated or consider using a [TST](#) approach where only animals in thin body condition are treated. If other options might exist, consider alternate species grazing, grazing summer annual forages, or hayfield grazing to create larvae-free, or reduced risk, places to graze because you will want these animals to reach breeding condition without setback.

Dry Ewes and Does



Fall:

What type of forage is being grazed?

Permanent Pasture

Hayfield

Dry Ewes and Does



Fall-permanent pasture:

Permanent pastures may have significant numbers of worm larvae on them if they were grazed during the spring and summer by lactating ewes or growing lambs and depending on the success of parasite control strategies. Dry ewes in good body condition are able to withstand a worm challenge that would create severe parasitism in lactating females or lambs. However, it is still a good idea to monitor the condition of these animals using the [FAMACHA©](#) system, body condition scoring, or [fecal egg counts](#), until the weather gets cold enough to retard transmission (ground temps below 50°F).



Fecal samples can be collected in resealable plastic bags.



The FAMACHA© system can be used to assess the overall situation of a group if records are kept.

Dry Ewes and Does



Fall-hayfield:

Hayfields not previously grazed this grazing season can provide a worm larvae-free place to graze in the fall and may provide excellent nutrition for flushing prior to breeding. [Strip grazing](#) with backfencing is the preferred technique. If alfalfa or clover is used, a bloat prevention strategy will need to be used. Care must be taken to allow the forage to recover before winter if the field is to be used for hay the following year.



Dry Ewes and Does



Winter:

Non-lactating mature females grazing in winter will be unlikely to acquire many worms if it is cold. Good options in some areas include standing corn, harvested corn fields, and stockpiled fescue.



Breeding Males



Breeding males.

Will these animals be grazing?

yes

no

Breeding males are usually managed as a separate group, and their nutrient requirements are principally for maintenance until breeding season. There is evidence that breeding males are somewhat more susceptible to severe parasitism than non-lactating adult females. This should be taken into consideration when these animals are grazed on a worm larvae-contaminated pasture, and a monitoring plan using [FAMACHA®](#), body condition scoring, or [fecal egg counting](#) should be implemented during the summer and fall grazing periods.

Return to [Table of Contents](#)116

Breeding Males



No grazing:

Mature rams and bucks kept indoors or in dry lot settings will get no exposure to [GIN](#).



Breeding Males



Yes-grazing season:

Season being grazed?

Spring

Summer

Fall

Winter

Breeding Males



Spring:

What type of forage is being grazed?

Permanent pasture

**Winter annual
(cereal rye, wheat)**

Hayfield

Breeding Males



Spring-permanent pasture:

When was the permanent pasture last grazed by sheep or goats?

Previous Spring

Previous Summer

Previous Fall

Previous Winter

Same Spring

Breeding Males



Spring-permanent pasture-previous spring/previous summer:

Pastures grazed in the spring that were last grazed the previous summer or spring should be relatively worm larvae-free. Worm larvae exposed to summer weather will use up their stored energy and be unlikely to survive winter. Breeding males should be able to withstand some parasite larval challenge with safety.



Breeding Males



Spring-permanent pasture-previous fall:

Although these pastures will likely have overwintered parasite larvae, breeding males that have had at least one season of grazing will have acquired some immunity to parasitism. However, research suggests that intact adult males may be more susceptible to parasitism than similar females. If the animals are in good body condition and pasture nutritive value is good, risk of severe parasitism is low. It would be a good idea to monitor body condition and eyelid score occasionally ([FAMACHA©](#) system).



More information about the FAMACHA© system can be found at:

<http://www.sheepandgoat.com/ACSRPC/Resources/famacha.html>

FAMACHA© cards and training may be available from your veterinarian or call your Extension educator to request training in your area.

STOP Return to “[Season being grazed](#)” 122

Breeding Males



Spring-permanent pasture-previous winter:

Worm eggs that may have been deposited on the pasture during winters where freezing temperatures are common do not survive well. The first two stages of larvae developing from eggs that do hatch during periods of warm weather in winter are very susceptible to drying. Third stage larvae are the infectious stage and survive much better. The risk of using pastures in spring that were last grazed during winter will depend on average temperatures, moisture, number of animals grazing, and their egg output. However, non-lactating animals can withstand some worm larval challenge without becoming severely parasitized. Use body condition score and/or use the [FAMACHA©](#) system to monitor these animals.

Breeding Males



Spring-permanent pasture-same spring:

Breeding males will be exposed to larvae developing from eggs shed in an earlier grazing this spring. If some strategy to minimize egg shedding during the earlier grazing is not employed, these pastures can have high levels of worm larvae. For example, if lactating females were the last to use this pasture, then deworming only thin ewes and ewes with twins and triplets can reduce pasture contamination while still providing a [refugia](#) of worms not exposed to dewormer.

Breeding Males



Spring-winter annual:

Winter annuals are usually planted after tillage of the soil and are usually planted in the fall following harvest of a crop. If they were planted on fields where sheep or goats had grazed that season, tillage will destroy most worm larvae making these pastures safe. If a “no-till” cultivation was used to plant the winter annual, some larvae may survive. No data exists to predict how many will survive, but it is likely to be only small numbers.



Breeding Males



Spring-hayfield:

If hayfields can be grazed in the spring, thus effectively harvesting “the first cutting” with the animals, they usually provide a worm larvae-free place to put them. Rams and bucks usually do not require a high level of nutrition at this time of year, and are usually not grazed on hayfields. If springtime hayfield grazing is anticipated, please see the information for [dry ewes/does](#).

Breeding Males



Summer:

What type of forage is being grazed?

Sericea lespedeza

Summer annual

Permanent pasture

Breeding Males



Summer: sericea lespedeza:

Information about planting and using sericea lespedeza can be found at the website for the [American Consortium for Small Ruminant Parasite Control](#).



STOP Return to "[Season being grazed](#)" 128

Breeding Males



Summer-summer annual:

Summer annuals such as turnips, kale, millet, cowpeas, or sorghum/sudangrass hybrids can provide a worm larvae-free place to graze mature breeding males. These forages are most effectively grazed using **strip grazing** with a back fence to prevent animals having access to previously grazed portions of the field. If the fences are moved across these clean pastures at 3-4 day intervals, and the back fence prevents animals from grazing where eggs may have been deposited, they will not acquire NEW infections as long as these larvae-free fields are available. Breeding males may not need the high quality nutrition that some summer annuals provide. Sorghum Sudangrass hybrids may be sufficient for breeding males. As with any other class of animals, avoid treating all the animals and immediately moving them to a worm larvae-free field as this will result in selection for dewormer resistant worms. If a **leader-follower** system is used where the males follow behind lactating females or weaned lambs/kids on these forages, the males may be exposed to high concentrations of parasite larvae unless the first animals to graze the forage have been managed to minimize their worm egg output.

Breeding Males



Summer-permanent pasture:

Permanent pastures grazed in summer can have very high levels of parasite larvae on them if they were grazed in the spring with lactating ewes or growing lambs that were shedding large numbers of parasite eggs. If only permanent pastures previously grazed by sheep or goats are available, consider using the [FAMACHA©](#) scoring system every 14-21 days throughout the summer to monitor. Alternatively, monitoring body condition and group level [fecal egg counts](#) may provide an alternative approach. If treatment is needed consider using the FAMACHA© system as a guide for which animals to be treated or consider using a [TST](#) approach where only animals in thin body condition are treated. If other options might exist, consider alternate species grazing, grazing summer annual forages, or hayfield grazing to create larvae-free, or reduced risk, places to graze because you will want these animals to reach breeding condition without setback.

Breeding Males



Fall:

What type of forage is being grazed?

Permanent Pasture

Hayfield

Breeding Males



Fall-permanent pasture:

Permanent pastures may have very high levels of parasite larvae on them if they were grazed in the summer with lactating females or growing lambs/kids that were shedding large numbers of parasite eggs. If only permanent pastures previously grazed by sheep or goats are available, consider using the [FAMACHA©](#) scoring system every 14-21 days throughout the summer to monitor parasitism. Monitoring body condition and group level [fecal egg counts](#) may provide an alternative approach. If treatment is needed consider using the FAMACHA© system as a guide for deciding which animals to treat or consider using a [TST](#) approach where only animals in thin body condition are treated. If other options might exist, consider alternate species grazing, grazing summer annual forages, or hayfield grazing to create larvae-free, or reduced risk places to graze.

Breeding Males



Fall-hayfield:

Hayfields not previously grazed this grazing season can provide a worm larvae-free place to graze in the fall and may provide excellent nutrition prior to breeding. [Strip grazing](#) with backfencing is the preferred technique. If alfalfa or clover is used, a bloat prevention strategy will need to be used. Care must be taken to allow the forage to recover before winter if the field is to be used for hay the following year.

Breeding Males



Winter:

Breeding rams grazing permanent pastures in winter will be unlikely to acquire many worms if it is cold. Good options in some areas include standing corn, harvested corn fields, and stockpiled fescue.



Stockpiled fescue in a field that has been set aside to grow from the end of summer through the fall.